



Clinical Report of Maximum Sexual

Conclusion on the treatment of 150 cases of impotence, premature ejaculation and menopause syndrome with Maximum sexual.

Insufficiency of both the spleen and kidney refers to a morbid condition mainly caused by failure of the insufficient kidney-yang to warm and nourish the spleen-yang, resulting in edema. It will be accompanied with the syndromes of lassitude, and so on. This is usually caused by insufficiency of yang, deficiency of kidney, excess of sexual intercourse, long-term illness, unclean diet, lassitude and tiredness, etc. 150 cases of impotence, premature ejaculation and menopausal syndrome were treated with Maximum sexual in our hospital. The clinical conclusion is as follows:

1. Case Source:

The cases of impotence and premature ejaculation are from the patient source of Out-patient Department, and the cases of menopausal syndrome are from the departments of Gynecology and In-patient.

2. Distribution of Age

Tablet 1 Distribution of Age among the Cases

Total Cases	Classification	Distribution of Age (Year)			
		18-22	23-30	31-40	41-50
150	Impotence (50)	8	7	15	20
	Premature Ejaculation(65)	25	14	16	10
	Menopausal syndrome (35)			7	28

3. Diagnosis Standard by Beijing University of Chinese Medicine and Pharmacology

3.1 Impotence

During the sexual intercourse, the penis of young man can't effectively erect and 75% of sexual opportunities can't be done. Then this is diagnosed as impotence in Beijing University of Chinese Medicine and Pharmacology

3.1.1 Classification of Impotence

Severe: Cannot have sexual intercourse for 3 month.



Medium: The successful rate of intercourse is less than 10% for 3 months.

Mild: The successful rate of intercourse is about 10-25% for 3 months.

3.1.2 Differentiation of Symptoms and Signs in Beijing University of Chinese Medicine and Pharmacology:

Decline of the Fire From the Gate of Life

Impotence, cold penis, aversion to cold, cold sperm, lingering diarrhea, pale, tongue fur, and weak pulse.

3.1.3 Diagnosis in Western Medicine

Diagnosis and Identification of Functional Impotence and Organic Impotence:

- A: Functional impotence occurs suddenly and organic impotence comes into being gradually except the ones caused by wounds or operation.
- B: Functional impotence may have an erectile phenomenon at night, but organic impotence does not.
- C: Functional impotence may have an erection under the stimulation of masturbation or erotic film, but organic impotence does not.

3.1.4 Standard of Curative Effectiveness

- A: Cured at sight: cured within 3 months, penis erects 90% and the successful rate of sexual intercourse reaches 70% or above.
- B: Remarkable Effectiveness: Erect to 90% after treatment, and the successful rate of sexual intercourse reaches 50% or above.
- C: Effectiveness: Erective function is improved after treatment, and the successful rate of sexual intercourse reaches 25% or above.
- D: No Effect: No improvement on erection before or after treatment.

3.2 Premature Ejaculation

3.2.1 Classification of Premature Ejaculation

- A: Mild: Penis inserts vagina and move for less than 1 minute, then ejaculates



B: Medium: once penis inserts vagina, it begins to ejaculate, or it ejaculates without body touch whenever the idea comes into his mind.

3.2.2 Standard of Curative Effectiveness

A: Cured at Sight: Sexual intercourse can be successfully done after 3 months treatment.

B: Remarkable Effectiveness: 75% of sexual intercourse is successful, and penis ejaculates 1 or more minutes later.

C: Effective: Penis can insert into vagina during the sexual intercourse, but it mostly ejaculates 1 minute later thrusting

D: No Effect: No change before or after treatment.

3.3 Menopausal Syndrome

A series of syndromes caused by decrease of hormone occur on woman before or after climacterium (menopausal period), including the syndrome of disorder of vegetative nervous function, are called menopausal syndrome.

3.3.1 Classification of Menopausal Syndrome

A: Mild: Hectic fever, sweat, facial flushing, and so on will last for several minutes each times or dozens of times each day. Unstable emotion, irrational, quarrelling and crying, hard to control oneself are the main symptoms and signs.

B: Medium: Dizziness, Headache, lassitude, tiredness, impression, insomnia and change of sexual desire.

C: Severe: Dizziness, tinnitus, short breath, palpitation, contraction of vulva and vagina, itching of vulva, pain and dryness of vagina, Contraction of uterus, dry skin, decrease of elasticity, contraction and descensus of breasts, etc.

3.3.2 Standard of Curative Effectiveness

A: Cured at Sight: All the clinical symptoms disappear after 3 months treatment.

B: Remarkable Effectiveness: Most of the clinical symptoms disappear after 3 months treatment.

C: Effective: The clinical symptoms decline and part of them disappear after 3 months



treatment.

D: No Effect: No big change before or after treatment.

3.3.3 Method of Treatment:

Oral administration of Maximum sexual, 2 tablets each time, 1 time each day, 1 month as a course of treatment, observation for another 2 months.

Results of Treatment:

A: Please see the Table 2 for the details of clinical treatment of 150 cases of impotence, premature ejaculation and menopausal syndrome

Table 2 Impotence, Premature Ejaculation and Menopausal Syndrome treated with Maximum sexual.

Total Curative Effectiveness Table

Group	Cases	Curative Effectiveness					
		Cured at Sight	Remarkable Effectiveness	Effective	No Effect	Remarkable Effective Rate	Total Effective Rate
Impotence	50	20	17	9	4	54%	92%
Premature	65	20	24	15	6	67.7%	90.8%
Menopausal Syndrome	35	8	14	8	5	62.9%	85.7%

B: Please see the Table 3 for the influence of Maximum sexual on the above-mentioned diseases in mild, medium and severe state

Tablet 3 Relationship Between Effectiveness and Diseases

Group	Type	Cases	Cured at Sight	Remarkable Effectiveness	Effective	No Effect	Total Effective Rate
Impotence	Mild	11	5	3	2	1	92%
	Medium	25	8	10	5	2	
	Severe	14	7	4	2	1	
Premature Ejaculation	Mild	15	4	6	5		90.8%
	Medium	29	10	11	6	2	
	Severe	21	6	7	4	4	
Menopausal Syndrome	Mild	14	4	6	2	2	85.7%
	Medium	13	3	5	3	2	
	Severe	8	2	3	2	1	

B: Maximum sexual can remarkably decrease the symptoms and signs of impotence, premature



ejaculation and menopausal syndrome, Most of patients have a remarkable curative effect after 1 month administration. Please see the following Table 4 for details.

Table 4 Improvement of Symptoms

Impotence, Premature Ejaculation				Menopausal Syndrome			
Symptoms and Signs Cases		Remarkable Effectiveness	Remarkable Effectiveness Rate %	Symptoms and Signs	Cases	Remarkable Effectiveness	Remarkable Effectiveness Rate %
Short Breath	95	70	73.7	Hectic Fever	30	23	76.7
Dizziness & Tinnitus	102	58	56.9	Sweat	25	18	72
Lassitude in Loin & Legs	114	95	83.3	Palpitation	32	25	78.1
Lassitude & Tiredness	110	82	74.5	Dizziness & Tinnitus	27	20	74.1
Impotence & Premature Ejaculation	106	60	56.6	Lassitude in Loin & Legs	34	29	85.3
Hyposexuality	88	70	79.5	Lassitude & Tiredness	28	21	75.0
Bronchitis due to Consumptive Diseases	62	48	77.4				

3.3.4 Results:

Maximum sexual is made from the natural Chinese herbs of Yohimbe bark extract, Cnidium Monnier, Epimedium extract, Songaria cynomorium, L-Arginine HCL Extract, Ginkgo Biloba Extract, Dahurian Angelica Root, Danshen Root, etc. it is specially used for hyposexuality, premature ejaculation, impotence, frigidity, and menopausal syndrome with the symptoms and signs of lassitude in loin and legs, aversion to cold, cold extremities (limbs), dizziness, tinnitus, short breath, palpitation, insomnia, lassitude and tiredness, etc. in the 150 cases of impotence, premature ejaculation and menopausal syndrome, the patients felt better and their sexual life was improved a lot after 1 month treatment with Maximum sexual. 2 month later, the total effective rate reached 92%. For the premature ejaculation patients, their successful intercourse rate has reached 75%, and they obtained a total curative effective rate of 90.8%. 20 out of them were cured during this period. As for menopausal syndrome, it is not easy to cure. After 2 months administration of Maximum sexual, the total effective rate reached 85.7%. This indicates that Maximum sexual has a very good curative effect on the above-mentioned diseases. No obvious toxic and side effect are found during the



clinical treatment.

6. Conclusion:

Maximum sexual is a kind of pure natural herbal preparation. In 150 cases of impotence, premature ejaculation and menopausal syndrome treated with Maximum sexual, the total curative effect reaches 85.7-92.0%. It has a much better effect on impotence and premature ejaculation. It can remarkably improve the treatment on the symptoms and signs of short breath, palpitation, dizziness, tinnitus, lassitude in loin and legs, lassitude and tiredness, impotence, premature ejaculation, hyposexuality and hectic fever. Maximum Sexual is safe and reliable, easy and convenient to carry.